arXiv:1304.3790v1 [math.AP] 13 Apr 2013

Global strong solution to Maxwell-Dirac equations in 1 + 1 dimensions

Aiguo You and Yongqian Zhang
School of Mathematical Sciences, Fudan University, Shanghai 200433
Key Laboratory of Mathematics for Nonlinear Sciences
Email address: yongqianz@fudan.edu.cn

April 16, 2013

Abstract

The Maxwell-Dirac equations with nonzero charge mass in one space dimension are studied under the Lorentz gauge condition. The global existence and uniqueness of solution in $C([0,+\infty); L^2(R^1)) \times C(R^1 \times [0,\infty))$ for initial value problem of Maxwell-Dirac equations are proved.

1 Introduction

We are concerned with the Maxwell-Dirac equations in one space dimension

$$\begin{cases} (i\gamma^\mu D_\mu + mI)\Psi = 0, \\ \partial_\mu F^{\mu\nu} = J^\nu, \end{cases} \tag{1.1}$$

under the Lorentz gauge condition

$$\partial_t A_0 - \partial_x A_1 = 0,$$

with

$$\begin{cases} \Psi_j(x,0) = \psi_j(x) \ \ (j = 1,2), \\ A_\nu(x,0) = a_\nu^0(x) \ \ (\nu = 0,1), \\ \partial_t A_\nu(x,0) = a_\nu^1(x) \ \ (\nu = 0,1), \end{cases} \tag{1.2}$$

which satisfy the constraint $a_0^1 - \partial_x a_1^0 = 0$. Here $D_\mu = \partial_\mu - iA_\mu$ is the covariant derivative and $F^{\mu\nu} = \partial_\nu A_\mu - \partial_\mu A_\nu$ is the curvature associated with the gauge field $A_\mu \in R$. Ψ denotes a two-spinor field defined on R^{1+1}, $J^\nu = \Psi^\dagger\gamma^0\gamma^\nu\Psi$ is a current density and $\Psi^\dagger = (\overline{\Psi_1}, \overline{\Psi_2})$ denotes the complex conjugate transpose of Ψ. $\partial_0 = \partial_t$, $\partial_1 = \partial_x$.

The Dirac gamma matrices are of the following:

$$\gamma^0 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \gamma^1 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

The system (1.1) can be rewritten as follows,

$$\begin{cases} \partial_t\Psi_1 + \partial_x\Psi_2 = im\Psi_1 + iA_0\Psi_1 + iA_1\Psi_2, \\ \partial_t\Psi_2 + \partial_x\Psi_1 = -im\Psi_2 + iA_0\Psi_2 + iA_1\Psi_1, \\ \Box A_0 = |\Psi_1|^2 + |\Psi_2|^2, \\ \Box A_1 = -(\overline{\Psi_2}\Psi_1 + \overline{\Psi_1}\Psi_2). \end{cases} \tag{1.3}$$

The global well-posedness of classical solution for the Maxwell-Dirac system in R^{1+1} has been established in [2]. Many works are devoted to study the solution in different functional spaces since then, see for instance [1]-[11] and the references therein. Recently in [10] Huh proved the global well-posedness of the strong solutions for the Maxwell-Dirac system(1.1) in R^{1+1}, where he assumed that the mass of charge is zero, that is, $m = 0$ in (1.1), and the solutions can be obtained by the explicit formula. For $m > 0$, as far as we know, there is no explicit formula for solutions. In this paper we consider more general case than that in [10], that is, $m \geq 0$. Our aim is to find the (Ψ, A) which solves the equation (1.3) with initial data (1.2) in the following sense and prove its uniqueness.

Definition 1.1 *(Ψ, A) with $\Psi \in L^2_{loc}(R^1 \times (0, +\infty))$ and $A \in C(R^1 \times [0, +\infty))$ is called a strong solution to (1.3) with the initial data (1.2) provided that there exists a sequence of smooth solutions $\{(\Psi^n, A^n_\mu)\}_{n=1}^\infty$ such that*

$$||\Psi^n - \Psi||_{L^2(K)} + ||A^n_1 - A_1||_{C(K)} + ||A^n_2 - A_2||_{C(K)} \to 0$$

as $n \to \infty$ for any compact set $K \subset R^1 \times [0, \infty)$, and

$$||\Psi^n(\cdot, 0) - \psi||_{L^2([a,b])} + ||A^n_\mu(\cdot, 0) - a^0_\mu||_{C([a,b])} + ||\partial_t A^n_\mu(\cdot, 0) - a^1_\mu||_{C([a,b])} \to 0$$

as $n \to \infty$ for $\mu = 1, 2$ and for any bounded interval $[a, b]$.

Remark 1.1 *For smooth solution to (1.3) with (1.2), which satisfies the constraint $a^1_0 - \partial_x a^0_1 = 0$, it was proved in [2] that*

$$\partial_t A_0 - \partial_x A_1 = 0.$$

Therefore, the Lorentz gauge condition also holds for the strong solution to (1.3) with (1.2). In the remaining, we only consider the Cauchy problem for (1.3).

The main result is presented as follows.

Theorem 1.1 *For the initial data $\psi = (\psi_1, \psi_2) \in L^2(R^1)$ and $a_\mu = (a_\mu^0, a_\mu^1) \in C(R^1)$, there exists a unique global strong solution (Ψ, A) to (1.3) with (1.2), which satisfy*

$$\Psi = (\Psi_1, \Psi_2) \in C([0,\infty); L^2(R^1)), A = (A_0, A_1) \in C(R^1 \times [0,\infty)).$$

The remaining of the paper is organized as follows. In section 2, we rewrite (1.1) in the equivalent form as (2.1-2.4). In section 3, based on the Chadam's result on the global H^1 strong solution for (1.1), we establish the key estimates in Lemmas 3.1 and 3.3 for classical solutions to (2.1-2.4), and get the uniform boundness on the solutions. In section 4, the key estimates given in Lemmas 3.1 and 3.3 play an important role in proving the precompactness of the approximate solutions $\{(u^n, v^n, A_\pm^n)\}_{n=1}^\infty$. Then we can get the convergence of the approximate solutions, this yields the existence of the global strong solution there. In section 5, we prove the uniqueness of the solutions.

2 Reduction of the problem

Denote

$$u = \Psi_1 + \Psi_2, v = \Psi_1 - \Psi_2,$$
$$A_+ = A_0 + A_1, A_- = A_0 - A_1,$$

and denote

$$u_0 = \psi_1 + \psi_2, v_0 = \psi_1 - \psi_2$$

and

$$a_\pm^0 = a_0^0 \pm a_1^0, a_\pm^1 = a_0^1 \pm a_1^1.$$

Then equations (1.3) can be written equivalently as

$$\partial_t u + \partial_x u = imv + iA_+ u, \tag{2.1}$$
$$\partial_t v - \partial_x v = imu + iA_- v, \tag{2.2}$$
$$\Box A_+ = |v|^2, \tag{2.3}$$
$$\Box A_- = |u|^2. \tag{2.4}$$

In addition, the initial data (1.2) can be written as

$$u(x,0) = u_0(x), \; v(x,0) = v_0(x) \tag{2.5}$$
$$A_\pm(x,0) = a_\pm^0(x), \; \frac{\partial A_\pm}{\partial t}(x,0) = a_\pm^1(x). \tag{2.6}$$

We will get the global solution to (1.3) with (1.2) by constructing the solutions to (2.1-2.6). That is, we will prove the following,

Theorem 2.1 *For any* $(u_0, v_0) \in L^2(R^1)$ *and* $(a^0_\pm, a^1_\pm) \in C(R^1)$*, there exists a unique strong solution* $(u, v, A_\pm)$ *to (2.1-2.6) with* $(u, v) \in C([0,\infty); L^2(R^1))$ *and* $A_\pm \in C(R^1 \times [0,\infty))$.

The strong solution to (2.1-2.4) is defined as in Definition 1.1.

3 Some Estimates on the smooth solutions

In this section, we assume that $\psi_1, \psi_2 \in C^\infty_c(R^1)$, $a^0_\nu \in C^\infty_c(R^1)$ and $a^1_\nu \in C^\infty_c(R^1)$. Due to the Chadam's result [2], the system (1.3) with initial data (1.2) has a unique solution $(\Psi, A) \in C^\infty$ with $supp\Psi(t,\cdot)$ and $suppA$ are compact sets in R for each $t \in R$. This implies that the problem (2.1-2.6) has a unique smooth solution $(u, v, A_\pm)$.

We will establish some estimates on the smooth solution $(u, v, A_\pm)$ in the next. To this end, we multiply (2.1) and (2.2) by $\overline{u}$ and $\overline{v}$ respectively, which yields the following

$$(|u|^2)_t + (|u|^2)_x = im(\overline{u}v - u\overline{v}), \tag{3.1}$$

$$(|v|^2)_t - (|v|^2)_x = im(\overline{v}u - v\overline{u}). \tag{3.2}$$

Then, direct computation shows that

Lemma 3.1 *Suppose* $\int_{-\infty}^{\infty}(|u_0(x)|^2 + |v_0(x)|^2)dx \le C_0$. *Then*

$$\int_{-\infty}^{\infty}(|u(x,t)|^2 + |v(x,t)|^2)dx = \int_{-\infty}^{\infty}(|u_0(x)|^2 + |v_0(x)|^2)dx \le C_0 \tag{3.3}$$

for $t \ge 0$.

Now, for any $(x_0, t_0) \in R^1 \times \overline{R_+}$, denote

$$\Delta(x_0, t_0) = \{(x,t) | t \in (0, t_0), x \in (x_0 - t_0 + t, x_0 + t_0 - t)\}$$

and

$$\Gamma_+(x_0, t_0) = \{(x,t) | t \in [0, t_0], x = x_0 - t_0 + t\},$$
$$\Gamma_-(x_0, t_0) = \{(x,t) | t \in [0, t_0], x = x_0 + t_0 - t\},$$

see Fig. 1.

Lemma 3.2 *For any* (x_0, t_0) *with* $t_0 > 0$*, there holds the following*

$$\int_{x_0-t_0+t}^{x_0+t_0-t}(|u(x,t)|^2 + |v(x,t)|^2)dx \le \int_{x_0-t_0}^{x_0+t_0}(|u_0(x)|^2 + |v_0(x)|^2)dx$$

for any $t \in [0, t_0]$.



Figure 1: Domain $\Delta(x_0, t_0)$

Proof. For $t \in (0, t_0)$, by (3.1-3.2) we have

$$(|u|^2 + |v|^2)_t + (|u|^2 - |v|^2)_x = 0,$$

then,

$$\begin{aligned}
\frac{d}{dt}\int_{x_0-t_0+t}^{x_0+t_0-t} (|u(x,t)|^2 + |v(x,t))|^2)dx &= \int_{x_0-t_0+t}^{x_0+t_0-t} \frac{\partial}{\partial t}(|u(x,t)|^2 + |v(x,t)|^2)dx \\
&- (|u(x_0+t_0-t,t)|^2 + |v(x_0+t_0-t,t)|^2) \\
&- (|u(x_0-t_0+t,t)|^2 + |v(x_0-t_0+t,t)|^2) \\
&= -2|u(x_0+t_0-t,t)|^2 - 2|v(x_0-t_0+t,t)|^2 \\
&\leq 0,
\end{aligned}$$

which gives the desired result. The proof is complete. □

Lemma 3.3 *There hold the following*

$$|u(x_0,t_0)|^2 \leq q(t_0)\int_{x_0-t_0}^{x_0+t_0} (|u_0(x)|^2 + |v_0(x)|^2)dx + e^{mt_0}|u_0(x_0-t_0)|^2,$$

$$|v(x_0,t_0)|^2 \leq q(t_0)\int_{x_0-t_0}^{x_0+t_0} (|u_0(x)|^2 + |v_0(x)|^2)dx + e^{mt_0}|v_0(x_0+t_0)|^2,$$

where $q(t_0) = e^{mt_0}m(mt_0+1)$.

Proof. Integrate the equation (3.1) along its characteristic curve $\Gamma_+(x_0, t_0)$, we have

$$\frac{d}{dt}|u(x_0 - t_0 + t, t)|^2 = im(\overline{u}v - \overline{v}u)(x_0 - t_0 + t, t).$$

Then

$$\frac{d}{dt}|u(x_0 - t_0 + t, t)|^2 \leq m(|u(x_0 - t_0 + t, t)|^2 + |v(x_0 - t_0 + t, t)|^2),$$

which leads to

$$|u(x_0 - t_0 + t, t)|^2 \leq e^{mt_0}(|u_0(x_0 - t_0)|^2 + m\int_{\Gamma_+} |v(x, t)|^2 ds). \tag{3.4}$$

To estimate the righthand side in the above, we integrate the equation (3.2) over the region $\Delta(x_0, t_0)$, then

$$|\int\int_{\Delta(x_0,t_0)} (|v|_t^2 - |v|_x^2) dxdt| = |im\int_0^{t_0}\int_{x_0-t_0+s}^{x_0+t_0-t} (\overline{v}u - \overline{u}v)(y, s)dyds|. \tag{3.5}$$

Applying the divergence theorem and Lemma 3.2 to the lefthand side and righthand side in (3.5) respectively, we have

$$|2\int_{\Gamma_+(x_0,t_0)} |v|^2 ds + \int_{x_0-t_0}^{x_0+t_0} |v_0|^2 dx| \leq mt_0(\int_{x_0-t_0}^{x_0+t_0} (|u_0|^2 + |v_0|^2)dx),$$

which leads to

$$\int_{\Gamma_+(x_0,t_0)} |v|^2 ds \leq (mt_0 + 1)(\int_{x_0-t_0}^{x_0+t_0} (|u_0|^2 + |v_0|^2)dx). \tag{3.6}$$

With (3.4) and (3.6), we have

$$|u(x_0 - t_0 + t, t)|^2 \leq e^{mt_0} m(mt_0 + 1)\int_{x_0-t_0}^{x_0+t_0} (|u_0|^2 + |v_0|^2)dx + e^{mt_0}|u_0(x_0 - t_0)|^2.$$

This gives the desired estimate for $|u|^2$. In the same way, we can get the estimate for $|v|^2$. The proof is complete. □

Lemma 3.4 *For any constant* $M \geq 0$,

$$\begin{aligned}\int_{|y|\geq M} |u(y, \tau)|^2 dy &\leq 2\tau q(\tau)\int_{|x|\geq M-\tau} (|u_0|^2 + |v_0|^2)dx \\ &\quad + e^{m\tau}\int_{|y|\geq M} |u_0(y - \tau)|^2 dy,\end{aligned}$$

$$\begin{aligned}\int_{|y|\geq M} |v(y, \tau)|^2 dy &\leq 2\tau q(\tau)\int_{|x|\geq M-\tau} (|u_0|^2 + |v_0|^2)dx \\ &\quad + e^{m\tau}\int_{|y|\geq M} |v_0(y + \tau)|^2 dy.\end{aligned}$$

Here $q(\tau) = e^{m\tau} m(m\tau + 1)$.

Proof. By Lemma 3.3, we have,

$$\begin{aligned}
\int_{|y|\geq M}|u(y,\tau)|^2dy &\leq \int_{|y|\geq M}q(\tau)\int_{y-\tau}^{y+\tau}(|u_0(x)|^2+|v_0(x)|^2)dxdy \\
&\quad+\int_{|y|\geq M}e^{m\tau}|u_0(y-\tau)|^2dy \\
&\leq e^{m\tau}\int_{|y|\geq M}|u_0(y-\tau)|^2dy \\
&\quad+q(\tau)\int_{|y|\geq M}\int_{-\tau}^{\tau}(|u_0(y+s)|^2+|v_0(y+s)|^2)dsdy \\
&\leq e^{m\tau}\int_{|y|\geq M}|u_0(y-\tau)|^2dy \\
&\quad+q(\tau)\int_{-\tau}^{\tau}\int_{|x-s|\geq M}(|u_0(x)|^2+|v_0(x)|^2)dxds \\
&\leq 2\tau q(\tau)\int_{|x|\geq M-\tau}(|u_0(x)|^2+|v_0(x)|^2)dx \\
&\quad+e^{m\tau}\int_{|y|\geq M}|u_0(y-\tau)|^2dy.
\end{aligned}$$

In the same way we can get the estimates on v. The proof is complete. □

For any (x_0,t_0) and (x_1,t_1) with $t_0\geq 0$ and $t_1\geq 0$, denote

$$\Omega(x_0,t_0;x_1,t_1)=\Big(\Delta(x_0,t_0)/\Delta(x_1,t_1)\Big)\cup\Big(\Delta(x_1,t_1)/\Delta(x_0,t_0)\Big),$$

and denote its Lesbeque measure by $meas(\Omega(x_0,t_0;x_1,t_1))$. It is obvious that

$$\lim_{(x,t)\to(x_0,t_0)} meas(\Omega(x_0,t_0;x,t))=0.$$

Lemma 3.5 *For any (x_0,t_0) and (x_1,t_1) with $t_0,t_1\in[0,T]$, there hold*

$$\begin{aligned}
&|\textstyle\int\int_{\Delta(x_0,t_0)}|u(y,\tau)|^2dyd\tau-\int\int_{\Delta(x_1,t_1)}|u(y,\tau)|^2dyd\tau| \\
&\leq C_0e^{mT}m(mT+1)meas(\Omega(x_0,t_0;x_1,t_1))+e^{mT}\textstyle\int\int_{\Omega(x_0,t_0;x_1,t_1)}|u_0(y-\tau)|^2dyd\tau,
\end{aligned}$$

and

$$\begin{aligned}
&|\textstyle\int\int_{\Delta(x_0,t_0)}|v(y,\tau)|^2dyd\tau-\int\int_{\Delta(x_1,t_1)}|v(y,\tau)|^2dyd\tau| \\
&\leq C_0e^{mT}m(mT+1)meas(\Omega(x_0,t_0;x_1,t_1))+e^{mT}\textstyle\int\int_{\Omega(x_0,t_0;x_1,t_1)}|v_0(y+\tau)|^2dyd\tau.
\end{aligned}$$

Proof. First we have

$$|\int\int_{\Delta(x_0,t_0)}|u(y,\tau)|^2dyd\tau-\int\int_{\Delta(x_1,t_1)}|u(y,\tau)|^2dyd\tau|=|\int\int_{\Omega(x_0,t_0;x_1,t_1)}|u(y,\tau)|^2dyd\tau|.$$

For the righthand side in above, it follows by Lemma 3.3 that

$$\begin{aligned}
\int\int_{\Omega(x_0,t_0;x_1,t_1)} |u(y,\tau)|^2 dyd\tau &\le \int\int_{\Omega(x_0,t_0;x_1,t_1)} e^{m\tau}|u_0(y-\tau)|^2 dyd\tau \\
&+ \int\int_{\Omega(x_0,t_0;x_1,t_1)} q(\tau)\int_{y-\tau}^{y+\tau}(|u_0|^2+|v_0|^2)dxdyd\tau \\
&\le e^{mT}\int\int_{\Omega(x_0,t_0;x_1,t_1)} |u_0(y-\tau)|^2 dyd\tau \\
&+ C_0 q(T) meas(\Omega(x_0,t_0;x_1,t_1)),
\end{aligned}$$

which proves the first inequality. In the same way we can prove the second inequality. The proof is complete. □

4 Existence of global strong solution

To get the strong solution to (2.1-2.6) under the assumption of Theorem 1.1, we choose sequences of smooth functions $u_0^n, v_0^n, a_{\pm n}^0, a_{\pm n}^1 \in C_c^\infty(R^1)$ such that the following hold.

(A1) There hold the following

$$\lim_{n\to\infty}\left(||u_0^n-u_0||_{L^2(R^1)}+||v_0^n-v_0||_{L^2(R^1)}\right)=0$$

and

$$\lim_{n\to\infty}\left(||a_{\pm n}^0-a_{\pm}^0||_{C([a,b])}+||a_{\pm n}^1-a_{\pm}^1||_{C([a,b])}\right)=0$$

for any bounded interval $[a,b]$.

(A2)

$$||u_0^n||_{L^2(R^1)}^2+||v_0^n||_{L^2(R^1)}^2\le C_0$$

and

$$\sup_{x\in R}|a_{\pm n}^j(x)|\le 2C_1$$

for $n\ge 1$ and $j=0,1$, where $C_1=\sup_{x\in R}|a_+^0(x)|+\sup_{x\in R}|a_+^1(x)|+\sup_{x\in R}|a_-^0(x)|+\sup_{x\in R}|a_-^1(x)|$.

Let (u^n,v^n,A_+^n,A_-^n) be the smooth solutions of (2.1-2.6) corresponding to the initial data $(u_0^n,v_0^n,a_{\pm n}^0,a_{\pm n}^1)$. Then, there hold the same estimates for (u^n,v^n,A_+^n,A_-^n) as in Section 3. Moreover,

$$2A_+^n(x,t)=a_{+n}^0(x+t)+a_{+n}^0(x-t)+\int_{x-t}^{x+t}a_{+n}^1(y)dy+\int_0^t\int_{x-t+s}^{x+t-s}|v^n(y,s)|^2dyds \tag{4.1}$$

and

$$2A_-^n(x,t) = a_{-n}^0(x+t)+a_{-n}^0(x-t)+\int_{x-t}^{x+t} a_{-n}^1(y)dy+\int_0^t \int_{x-t+s}^{x+t-s} |u^n(y,s)|^2 dyds. \tag{4.2}$$

Lemma 4.1 *For any $T \geq 0$, $n \geq 1$, there holds*

$$\sup_{n\geq 0} \sup_{x\in R^1, t\in[0,T]} |A_\pm^n(x,t)| \leq C_1(T+1)+C_0T.$$

Proof. By Lemma 3.1, we can deduce the result from (4.1-4.2). The proof is complete. □

Lemma 4.2 *There are functions $A_\pm^* \in C(R^1\times[0,\infty))$ and subsequences of $\{A_\pm^n\}_{n=1}^\infty$, still denoted by $\{A_\pm^n\}_{n=1}^\infty$, such that*

$$||A_+^n - A_+^*||_{C(K)} + ||A_-^n - A_-^*||_{C(K)} \to 0$$

as $n\to\infty$ for any compact set $K \subset R^1\times[0,\infty)$.

Proof. We consider the sequence $\{A_+^n\}_{n=1}^\infty$. Due to the Lemma 4.1, it suffices to prove that $\{\int_0^t\int_{x-t+s}^{x+t-s}|v^n(y,s)|^2dyds\}_{n=1}^\infty$ is equicontinuous in any domain $\Delta(X,T)$ with $T\geq 0$.

By Lemma 3.5, for any (x_0,t_0) and (x_1,t_1) with $t_0,t_1\in[0,T]$, we have

$$\begin{aligned} &|\textstyle\int\int_{\Delta(x_0,t_0)}|v^n(y,\tau)|^2dyd\tau - \int\int_{\Delta(x_1,t_1)}|v^n(y,\tau)|^2dyd\tau| \\ &\leq C(T)meas(\Omega(x_0,t_0;x_1,t_1)) + C(T)\textstyle\int\int_{\Omega(x_0,t_0;x_1,t_1)}|v_0^n(y+\tau)|^2dyd\tau, \\ &\leq C(T)meas(\Omega(x_0,t_0;x_1,t_1)) + C(T)\textstyle\int\int_{\Omega(x_0,t_0;x_1,t_1)}|v_0(y+\tau)|^2dyd\tau \\ &\qquad +TC(T)||v_0^n - v_0||_{L^2(R^1)}^2, \end{aligned} \tag{4.3}$$

where $C(T)=e^{mT}(m(mT+1)+1)$. Then, by the continuity of the function $\int\int_{\Omega(x_0,t_0;x_1,t_1)}|v_0(y+\tau)|^2dyd\tau$ and the convergence of the sequence $\{v^n\}_{n=1}^\infty$, it follows from (4.3) that $\{\int_0^t\int_{x-t+s}^{x+t-s}|v^n(y,s)|^2dyds\}_{n=1}^\infty$ is equicontinuous in the domain $\Delta(X,T)$. Therefore the collection $\{A_+^n\}_{n=1}^\infty$ is equicontinuous in the domain $\Delta(X,T)$.

In the same way, we can prove that $\{A_-^n\}_{n=1}^\infty$ is equicontinuous in the domain $\Delta(X,T)$. Then, by Arzela-Ascoli Theorem ([12], pp 245), we can have the desired result. The proof is complete. □

Lemma 4.3 *Let $\{A_\pm^n\}$ be the sequences given by Lemma 4.2, For any $T\geq 0$,*

$$\lim_{k,l\to\infty}\int_0^T\int_{R^1}(|A_+^k - A_+^l|^2 + |A_-^k - A_-^l|^2)(|u^k|^2+|v^k|^2)(x,\tau)dxd\tau = 0.$$

Proof. For any $M > 0$, denote

$$I_{M,1}^{k,l} = \int_0^T \int_{|x|\geq M} (|A_+^k - A_+^l|^2 + |A_-^k - A_-^l|^2)(|u^k|^2 + |v^k|^2)(x,\tau)dxd\tau$$

and

$$I_{M,2}^{k,l} = \int_0^T \int_{|x|\leq M} (|A_+^k - A_+^l|^2 + |A_-^k - A_-^l|^2)(|u^k|^2 + |v^k|^2)(x,\tau)dxd\tau.$$

By Lemmas 4.1, 3.1 and 3.5, we have

$$\begin{aligned} I_{M,1}^{k,l} &\leq (C_1(T+1) + C_0T) \int_0^T \int_{|x|\geq M} (|u^k|^2 + |v^k|^2)(x,\tau)dxd\tau \\ &\leq C_T'\Big(T \int_{|x|\geq M-T} (|u_0^k|^2 + |v_0^k|^2)dx + \int_0^T \int_{|y|\geq M} |u_0^k(y-\tau)|^2 dyd\tau \\ &\quad + \int_0^T \int_{|y|\geq M} |v_0^k(y+\tau)|^2 dyd\tau\Big) \\ &\leq C_T'\Big(2T \int_{|x|\geq M-T} (|u_0^k|^2 + |v_0^k|^2)dx\Big) \\ &\leq 2TC_T'\Big(\int_{|x|\geq M-T} (|u_0|^2 + |v_0|^2)dx + ||u_0 - u_0^k||_{L^2}^2 + ||v_0 - v_0^k||_{L^2}^2\Big) \end{aligned}$$

and

$$I_{M,2}^{k,l} \leq C_0(||A_+^k - A_+^l||_{C(K)}^2 + ||A_-^k - A_-^l||_{C(K)}^2),$$

where $K = [-M, M] \times [0,T]$ and $C_T' = 2e^{mT}(m(mT+1)+1)(C_1(T+1)+C_0T)$. Then, by Lemma 4.2, we have

$$\begin{aligned} &\limsup_{k,l\to\infty} \int_0^T \int_R (|A_+^k - A_+^l|^2 + |A_-^k - A_-^l|^2)(|u^k|^2 + |v^k|^2)(x,\tau)dxd\tau \\ &\leq \limsup_{k,l\to\infty} (I_{M,1}^{k,l} + I_{M,2}^{k,l}) \\ &\leq 2TC_T'\Big(\int_{|x|\geq M-T} (|u_0|^2 + |v_0|^2)dx\Big), \end{aligned}$$

which implies the desired result by letting $M \to \infty$. The proof is complete. □

Proof of the existence of solutions in Theorem 2.1. Let $\{A_\pm^n\}$ be the sequences given by Lemma 4.2. Due to Lemma 4.2, it suffices to prove the convergence of $\{(u^k, v^k)\}_{k=1}^\infty \in (R^1 \times (0,T))$, for any $T \geq 0$.

With $(u^n, v^n, A_\pm^n)$ of $(2.1) - (2.4)$, we deduce that

$$\begin{aligned} (|u^k - u^l|^2)_t + (|u^k - u^l|^2)_x &= im((\overline{u}^k - \overline{u}^l)(v^k - v^l) - (u^k - u^l)(\overline{v}^k - \overline{v}^l)) + \\ &\quad i(A_+^k - A_+^l)u^k(\overline{u}^k - \overline{u}^l) - i(A_+^k - A_+^l)\overline{u}^k(u^k - u^l), \\ (|v^k - v^l|^2)_t - (|v^k - v^l|^2)_x &= -im((\overline{u}^k - \overline{u}^l)(v^k - v^l) - (u^k - u^l)(\overline{v}^k - \overline{v}^l)) + \\ &\quad i(A_-^k - A_-^l)v^k(\overline{v}^k - \overline{v}^l) - i(A_-^k - A_-^l)\overline{v}^k(v^k - v^l). \end{aligned}$$

Then, there exists a constant C_2 independent of k, l such that

$$\frac{d}{dt}\int_R (|u^k - u^l|^2 + |v^k - v^l|^2)dx \le C_2 \int_R (|u^k - u^l|^2 + |v^k - v^l|^2)dx$$
$$+C_2 \int (|A_+^k - A_+^l|^2 + |A_-^k - A_-^l|^2)(|u^k|^2 + |v^k|^2)dx$$

for $t \ge 0$. Applying the Gronwall inequality to this inequality gives

$$\int_R (|u^k - u^l|^2 + |v^k - v^l|^2)(x,t)dx \le e^{C_2 t}\Big(||u_0^k - u_0^l||^2_{L^2(R^1)} + ||v_0^v - u_0^l||^2_{L^2(R^1)}$$
$$+C_2 e^{C_2 t}\int_0^t \int_R (|A_+^k - A_+^l|^2 + |A_-^k - A_-^l|^2)(|u^k|^2 + |v^k|^2)(x,\tau)dxd\tau\Big)$$

for $t \ge 0$, which implies $\{(u^k, v^k)\}_{k=1}^{\infty}$ is a Cauchy sequence in $C([0,T], L^2(R^1))$ by Lemma 4.3 for any $T \ge 0$. Therefore, there exists a function $(u^*, v^*) \in C([0,T], L^2(R^1))$ such that

$$||u^k - u^*||_{C([0,T],L^2(R^1))} + ||v^k - v^*||_{C([0,T],L^2(R^1))} \to 0$$

as $k \to \infty$ for any $T \ge 0$. Thus this gives the proof of existence of solution. We will prove the uniqueness in the next section. $\square$

5 Uniqueness of global solutions

In this section, we prove the uniqueness of the strong solution which we constructed above.

Let $(u_1, v_1, A_\pm)$ and $(u_2, v_2, B_\pm)$ be two strong solutions with the same initial data, that is, $u_1(x,0) = u_2(x,0)$, $v_1(x,0) = v_2(x,0)$, $A_\pm(x,0) = B_\pm(x,0)$ and $\partial_t A_\pm(x,0) = \partial_t B_\pm(x,0)$. Our goal is to prove that $u_1 = u_2$, $v_1 = v_2$, $A_\pm = B_\pm$. We assume that $(u_1^n, v_1^n, A_\pm^n)$ and $(u_2^n, v_2^n, B_\pm^n)$ are two sequences of smooth solutions of equations (2.1-2.4), such that $(u_1^n, v_1^n, A_\pm^n) \to (u_1, v_1, A_\pm)$, $(u_2^n, v_2^n, B_\pm^n) \to (u_2, v_2, B_\pm)$ in the sense of Definition 1.1.

Denote $w_1^n = u_1^n - u_2^n$, $w_2^n = v_1^n - v_2^n$ and $G_\pm^n = A_\pm^n - B_\pm^n$, then

$$\begin{aligned} \partial_t w_1^n + \partial_x w_1^n &= imw_2^n + G_+^n u_1^n + iB_+^n w_1^n, \\ \partial_t w_2^n - \partial_x w_2^n &= imw_1^n + G_-^n v_1^n + iB_-^n w_2^n, \\ \Box G_+^n &= \overline{v_1^n} w_2^n + \overline{w_2^n} v_2^n, \\ \Box G_-^n &= \overline{u_1^n} w_1^n + \overline{w_1^n} u_2^n, \end{aligned}$$

which gives the following,

$$\partial_t |w_1^n|^2 + \partial_x |w_1^n|^2 = im(w_2^n \overline{w_1^n} - w_1^n \overline{w_2^n}) + 2Im(G_+^n \overline{u_1^n} w_1^n), \tag{5.1}$$
$$\partial_t |w_2^n|^2 - \partial_x |w_2^n|^2 = im(w_1^n \overline{w_2^n} - w_2^n \overline{w_1^n}) + 2Im(G_-^n \overline{v_1^n} w_2^n), \tag{5.2}$$
$$\Box G_+^n = \overline{v_1^n} w_2^n + \overline{w_2^n} v_2^n, \tag{5.3}$$
$$\Box G_-^n = \overline{u_1^n} w_1^n + \overline{w_1^n} u_2^n. \tag{5.4}$$

By (5.1) and (5.2), we have

$$\partial_t(|w_1^n|^2+|w_2^n|^2)+\partial_x(|w_1^n|^2-|w_2^n|^2)=2Im(G_+^n\overline{u_1^n}w_1^n)+2Im(G_-^n\overline{v_1^n}w_2^n),$$

for $0\le t\le T$.

Integrating the above equation over the interval $[-T+t,T-t]$, we have

$$\begin{aligned}\frac{d}{dt}(\int_{-T+t}^{T-t}(|w_1^n(x,t)|^2+|w_2^n(x,t)|^2)dx) &+ 2|w_1^n(T-t,t)|^2+2|w_2^n(-T+t,t)|^2\\ &\le 2\int_{-T+t}^{T-t}(|G_+^n\overline{u_1^n}w_1^n|+G_-^n\overline{v_1^n}w_2^n|)dx.\end{aligned}$$

Denote $I^n(t,T)=\int_{-T+t}^{T-t}(|w_1^n(x,t)|^2+|w_2^n(x,t)|^2)dx$. Then,

$$\begin{aligned}\frac{d}{dt}I^n(t,T) &\le 2||G_+^n||_{C(D_T)}||u_1^n||_{L^2(D_T)}||w_1^n||_{L^2(D_T)}\\ &+ 2||G_-^n||_{C(D_T)}||v_1^n||_{L^2(D_T)}||w_2^n||_{L^2(D_T)}, \qquad (5.5)\end{aligned}$$

where $D_T=\{(x,t)|-T+t<x<T-t,0\le t\le T\}$.

Applying the D'Alembert formula to the equations (5.3-5.4), we have,

$$\begin{aligned}2G_+^n &= g_{0+}^n(x-t)+g_{0+}^n(x+t)+\int_{x-t}^{x+t}g_{1+}^n(\tau)d\tau\\ &+\int_0^t\int_{x-t+s}^{x+t-s}(\overline{v_1^n}w_2^n+\overline{w_2^n}v_2^n)(y,s)dyds,\\ 2G_-^n &= g_{0-}^n(x-t)+g_{0-}^n(x+t)+\int_{x-t}^{x+t}g_{1-}^n(\tau)d\tau\\ &\int_0^t\int_{x-t+s}^{x+t-s}(\overline{u_1^n}w_1^n+\overline{w_1^n}u_2^n)(y,s)dyds,\end{aligned}$$

where $g_{0\pm}^n(x)=G_\pm^n(x,0)$ and $g_{1\pm}^n(x)=\frac{\partial G_\pm^n}{\partial t}(x,0)$.

Then,

$$\begin{aligned}||G_+^n||_{C(D_T)} &\le ||g_{0+}^n||_{C([-T,T])}+T||g_{1+}^n||_{(C[-T,T])}\\ &+\frac{1}{2}(||v_1^n||_{L^2(D_T)}+||v_2^n||_{L^2(D_T)})||w_2^n||_{L^2(D_T)}, \qquad (5.6)\\ ||G_-^n||_{C(D_T)} &\le ||g_{0-}^n||_{C([-T,T])}+T||g_{1-}^n||_{C([-T,T])}\\ &+\frac{1}{2}(||u_1^n||_{L^2(D_T)}+||u_2^n||_{L^2(D_T)})||w_1^n||_{L^2(D_T)}. \qquad (5.7)\end{aligned}$$

With (5.5),(5.6) and (5.7), we have

$$\frac{d}{dt}(I^n(t,T))\le C_1(I^n(t,T))+C_2g^n(T)$$

for some constants $C_1 > 0$ and $C_2 > 0$ depending only on C_0, where Lemma 3.1 is used. Here $g^n(T) = ||g^n_{0+}||^2_{C([-T,T])} + T^2||g^n_{1+}||^2_{(C[-T,T])} + ||g^n_{0-}||^2_{C([-T,T])} + T^2||g^n_{1-}||^2_{C([-T,T])}$.

Then applying the Gronwall inequality and lemma 3.1 to above, we can get

$$I^n(t,T) \leq (I^n(0,T) + \frac{C_2}{C_1} g^n(T))e^{C_1 t},$$

for $t \geq 0$ and $T \geq 0$. Due to the Definition 1.1,

$$\lim_{n\to 0} I^n(0,T) = 0, \quad \lim_{n\to 0} g^n(T) = 0,$$

which lead to

$$\begin{aligned} \int_{-T+t}^{T-t} (|u_1(y,t) - u_2(y,t)|^2 + |v_1(y,t) - v_2(y,t)|^2) dy &= \lim_{n\to\infty} I^n(t,T) \\ &= 0. \end{aligned}$$

Then $u_1 \equiv v_1, u_2 \equiv v_2$.

Now by (5.6), (5.7), we have

$$\begin{aligned} ||A_+ - B_+||_{C(D_T)} &= \lim_{n\to\infty} ||G^n_+||_{C(D_T)} = 0, \\ ||A_- - B_-||_{C(D_T)} &= \lim_{n\to\infty} ||G^n_-||_{C(D_T)} = 0, \end{aligned}$$

therefore $A_\pm = B_\pm$. The proof is complete.□

Acknowledgement

This work was partially supported by NSFC Project 11031001 and 11121101, by the 111 Project B08018 and by Ministry of Education of China.

References

[1] N. Bournaveas, *Local existence for the Maxwell-Dirac equations in three space dimensions,* Comm. Partial Diff. Eqs 21(1996), 693-720.

[2] J. Chadam, *Global solution of the Cauchy problem for the (classical) coupled Maxwell-Dirac equations in one space dimension,* J. Funct. Anal. 13(1973), 173-184.

[3] J. Chadam and R. Glassey, *On the Maxwell-Dirac equations with zero magnetic field and their solution in two space dimensions,* J. Math. Anal. Appl. 53(1976), 495-507.

[4] P. D'Ancona, D. Foschi and S. Selberg, *Null structure and almost optimal local well-posedness of the Maxwell- Dirac system,* Amer. J. Math. 132 (2010), no. 3, 771-839.

[5] P. D'Ancona and S. Selberg, *Global well-posedness of the Maxwell-Dirac system in two space dimensions,* J. Funct. Anal. 260 (2011), no. 8, 2300-2365.

[6] V. Delgado, *Global solutions of the Cauchy problem for the (classical) coupled Maxwell-Dirac and other nonlinear Dirac equations in one space dimension,* proc. Am. Math. Soc. 69(1978), 289-296.

[7] M. Flato, J. Simon and E. Taflin, *Asymptotic sompleteness, global existence and the infrared problem for the Maxwell-Dirac equations,* Memoirs Amer. Math. Soc. 127(1997), no.606.

[8] L. Gross, *The Cauchy problem for the coupled Maxwell and Dirac equations,* Comm. Pure. Appl. Math. 19(1996), 1-15.

[9] V. Georgiev,*Small amplitude solutions of the Maxwell-Dirac equations* Indiana Univ. Math. J. 40 (1991), no. 3, 845-883.

[10] H. Huh, *Global charge solutions of Maxwell-Dirac equations in* R^{1+1}, J. Phys, A: Math. Theor. 43 (2010), 445206 (7pp).

[11] M. Okamoto, *Well-posedness and ill-posedness of the Cauchy problem for the Maxwell-Dirac system in 1+1 space time dimensions,* Adv. Differential Equations 18 (2013) no. 1/2, 179-199.

[12] W. Rudin, *Real and Complex Analysis,* McGraw-Hill Book Company, 1987.

[13] Y. Zhou, *Uniqueness of weak solutions of 1+1 dimensional wave maps,* Math. Z. 232 (1999), no. 4, 707-719.